

GRAVITY & DUALITY

Supersymmetric 6-D gravity with (4,0) Susy

Schloss Rindberg, November 2016



Ioannis Bakas
1960-2016

Theory X?

- Considerable evidence for mysterious interacting 6-D (2,0) non-lagrangian SCFT
- Key to understanding SYM in $D < 6$, S-duality
- Similar story for gravity?
- IF there is an interacting (4,0) SCFT in 6-D, it would be exotic CONFORMAL theory giving SUGRA in $D < 6$

(2,0) Theory

- Free (2,0) theory in 6-D: 2-form B , $H=H^*$
- Reduces to 5-D $N=4$ Maxwell, $F=dA$
- Interacting (2,0) SCFT, non-lagrangian, reduces to 5-D SYM
- Strong coupling limit of 5-D SYM: (2,0) SCFT
- Stringy constructions: M5-brane, IIB on $K3$

(4,0) Theory


- Free (4,0) theory in 6-D: SCFT
- Reduces to 5-D linearised N=8 SUGRA
- Is there an interacting (4,0) SCFT? Non-lagrangian, reducing to 5-D SUGRA?
- Strong coupling limit of 5-D SUGRA?
- Exotic conformal theory of gravity?
- Highly symmetric (4,0) phase of M-theory?

Gravity = (YM)²

- Free SUGRA ~ Free (SYM)²



- Free (4,0) ~ Free ((2,0) theory)²



- Free (2,0) reduces to 5-D theory of photon + dual photon
- Free (4,0) reduces to 5-D theory of graviton + dual graviton + double dual graviton

5-D Superalgebra

$$\{Q_\alpha^a, Q_\beta^b\} = \Omega^{ab} (\Gamma^\mu C)_{\alpha\beta} P_\mu + C_{\alpha\beta} (Z^{ab} + \Omega^{ab} K)$$

- Central charges Z, K
- Z Electric charges for Maxwell fields
- States with K ~ KK modes of 6-D (p,0) theory
- SYM: K carried by BPS solitons (from YM instantons)
- Does M-theory on T^6 have BPS states with K?
- Do they become massless at strong coupling?


Maxwell in D- dimensions

- Photon A_μ
- Dual photon: $n=D-3$ form $\tilde{A}_{\mu_1 \dots \mu_n}$
 $F = *\tilde{F}$
- Magnetic charges: D-4 branes.
A has Dirac strings, or connection on non-trivial bundle, \tilde{A} well-defined
- Electric charges: 0-branes.
 \tilde{A} has Dirac string singularities, A OK
- YM? No non-abelian theory for \tilde{A}




Linearised Gravity

- **Graviton** $h_{\mu\nu}$  $(1, 1)$

Field strength $R_{\mu\nu\rho\sigma}$  $(2, 2)$

- **Dual Graviton**


$\tilde{h}_{\mu_1 \dots \mu_n \nu}$ $(n, 1)$



$n=D-3$


- **Double Dual Graviton**

$\tilde{\tilde{h}}_{\mu_1 \dots \mu_n \nu_1 \dots \nu_n}$
 (n, n)




Linearised Gravity

- **Graviton** $h_{\mu\nu}$  $(1, 1)$

Field strength $R_{\mu\nu\rho\sigma}$  $(2, 2)$

- **Dual Graviton**

$\tilde{h}_{\mu_1 \dots \mu_n \nu}$ $(n, 1)$



$\tilde{R}_{\mu_1 \dots \mu_{n+1} \rho\sigma}$


$n=D-3$

$(n + 1, 2)$

- **Double Dual Graviton**

$\tilde{\tilde{h}}_{\mu_1 \dots \mu_n \nu_1 \dots \nu_n}$

(n, n)



$\tilde{\tilde{R}}_{\mu_1 \dots \mu_{n+1} \nu_1 \dots \nu_{n+1}}$

$(n + 1, n + 1)$

Field strengths are Dual:

$$R \quad \tilde{R} = *R \quad \tilde{\tilde{R}} = *R*$$

Duality Exchanges field equals and Bianchis

$$R_{\mu\rho\nu}{}^\rho = 0 \quad \leftrightarrow \quad \tilde{R}_{[\mu_1 \dots \mu_n \mu_{n+1} \nu]\rho} = 0$$

$$R_{[\mu\nu\rho]\sigma} = 0 \quad \leftrightarrow \quad \tilde{R}_{\mu_1 \dots \mu_n \rho \nu}{}^\rho = 0$$

Electric and Magnetic Grav Sources T, \tilde{T} for h, \tilde{h}

$$\begin{array}{ll} \tilde{T} : & \text{Dirac strings for } h \\ T : & \text{Dirac strings for } \tilde{h} \end{array}$$

- **Hull 2000**: Dual graviton, double dual graviton in D dims, motivated by 6-D CFT
- **West 2001**: Dual graviton & E_{11}
- **Bekaert, Boulanger & Henneaux 2002**: No interactions for dual graviton, no dual formulation of GR
- Non-linear action with both
West 2001, Boulanger & Hohm 2008
 $D=11$ SUGRA: Bergshoeff, de Roo & Hohm

D=6 (2,0) free theory

R-symmetry $Sp(2)=USp(4)$

Superconformal $OSp(4/8^*) \supset USp(4) \times SO(6,2)$

$$B_{MN} \quad H = *H$$

5 scalars, 4 fermions

Reduce to D=5

$$B_{\mu\nu}, B_{\mu 5} = A_{\mu} \quad H = *F$$

A,B dual, not independent

A, 5 scalars, 4 fermions: **D=5 N=4 vector multiplet**

Reduce to D=4

2 vector fields $B_{\mu i} = A_{\mu i} \quad i = 1, 2 \quad F_1 = *F_2$

SL(2,Z): diffeos on T^2 (A_1, A_2) doublet

Only one independent field, **D=4 N=4 vector multiplet**

SL(2,Z): (A_1, \tilde{A}_1) doublet, E-M duality

The (4,0) Supermultiplet

D=6 little group $SO(4) \sim SU(2) \times SU(2)$

States in representations of $SU(2) \times SU(2) \times USp(8)$

$$(5, 1; 1) + (4, 1; 8) + (3, 1; 27) + (2, 1; 48) + (1, 1; 42)$$

Light-cone fields $\alpha, \beta = 1, 2 \quad a, b = 1, \dots, 8$

$$C_{\alpha\beta\gamma\delta}, \psi_{\alpha\beta\gamma}^a, B_{\alpha\beta}^{ab}, \lambda_{\alpha}^{abc}, \phi^{abcd}$$

Covariant fields

$$C_{MNPQ}, \psi_{MN}^a, B_{MN}^{ab}, \lambda^{abc}, \phi^{abcd}$$

D=6 Free (4,0) Theory


Hull

42 scalars

27 self-dual B_2 : $H = *H$


Gauge field

C_{MNPQ}



Curvature

G_{MNPQRS}



Self-dual: $G = *G = G*$

“Supergravity without a graviton”

Superconformal $OSp(8/8^*) \supset USp(8) \times SO(6,2)$

Reduce to D=5

27 $B_2 \rightarrow 27$ vectors A_1 , 42 scalars $\rightarrow 42$ scalars

$$C_{\mu 5 \nu 5} = h_{\mu \nu}$$

$$C_{\mu \nu \rho 5} = \tilde{h}_{\mu \nu \rho}$$

$$C_{\mu \nu \rho \sigma} = \tilde{\tilde{h}}_{\mu \nu \rho \sigma}$$

Self-duality: Only one of these independent, dual gravitons


Spectrum of D=5 N=8 SUGRA!

Graviton, 27 vectors, 42 scalars Diffeos

Vectors from B_{MN}

Graviton from C_{MNPQ}

Diffeos from C gauge transformations. Parameter



Reduce to D=4

42 scalars \rightarrow 42 scalars, Dual vector doublets $B_{\mu i} = A_{\mu i}$

Metrics $C_{\mu(ij)\nu} = -(h_{\mu\nu})_{ij}$

Curvatures: $R_{21} = *R_{11}$, $R_{12} = R_{11}^*$, $R_{22} = *R_{11}^*$

$$h_{21} = \tilde{h}_{11}, \quad h_{22} = \tilde{\tilde{h}}_{11}$$

Just h_{11} independent

SL(2,Z) on torus:

(A_1, A_2) doublets, E-M duality

Triplet h_{ij} : gravitational triality symmetry of free theory

5-D SYM at Strong Coupling

$$\{Q_\alpha^a, Q_\beta^b\} = \Omega^{ab} (\Gamma^\mu C)_{\alpha\beta} P_\mu + C_{\alpha\beta} (Z^{ab} + \Omega^{ab} K)$$

Z electric charges: carried by W-bosons etc

YM instanton in R^4 lifts to BPS soliton in 5-D

K proportional to instanton number n , (2,0) short mult.

$$M \propto \frac{n}{g_{YM}^2}$$

Light at strong coupling: KK tower for 6'th dimension

Decompactifies to (2,0) theory in 6D as $g_{YM}^2 \rightarrow \infty$

(2,0) Interacting CFT

D=5 non-renormalizable, defined within string theory
e.g. D4 brane theory

Strong coupling limit defined within string theory
e.g. multiple D4 branes \rightarrow multiple M5 branes

No direct construction of interacting (2,0) theory.

Reduce on T^2 gives interacting N=4 SYM
and $SL(2, \mathbb{Z})$ S-duality from torus diffeos

g_{YM} dimensionful. Limit is one to high energies

$$E \gg (g_{YM})^{-2}$$

$$E(g_{YM})^2 \rightarrow \infty$$

SUGRA at Strong Coupling

$$\{Q_\alpha^a, Q_\beta^b\} = \Omega^{ab} (\Gamma^\mu C)_{\alpha\beta} P_\mu + C_{\alpha\beta} (Z^{ab} + \Omega^{ab} K)$$

If there are BPS states carrying K , with spectrum

$$M \propto \frac{n}{l_{Plank}}$$

Become light in strong coupling (high energy) limit

$$E \times l_{Plank} \rightarrow \infty$$

Decompactification limit with K -states as a KK tower?
If so, must decompactify to a (4,0) theory in 6D as (4,0)
short multiplet

M-Theory Compactified on a Torus

D=4:

28 vector fields

28 electric + 28 magnetic charges

$E_7(\mathbb{Z})$ symmetry

D=5:

27 vector fields

27 electric charges Z^{ab} + 27 magnetic strings

$E_6(\mathbb{Z})$ symmetry

“Topological” charge K , carried by KK monopoles

Reduce $5 \rightarrow 4$:

Graviphoton $g_{\mu 5}$ Electric charge: P^5 Magnetic charge: K

D=5 N=8 Superalgebra

$$\{Q_\alpha^a, Q_\beta^b\} = \Omega^{ab} (\Gamma^\mu C)_{\alpha\beta} P_\mu + C_{\alpha\beta} (Z^{ab} + \Omega^{ab} K)$$

K carried by KK monopoles

Gibbons & Perry

Z^{ab} carried by charged 0-branes (from wrapped M-branes)

BPS bound

$$M \geq |K|$$

Full D=5 M-theory on S^1 :

No killing vectors, full KK tower etc

Has $E_7(Z)$ symmetry

Includes duality $P^5 \leftrightarrow K$

D>5: D-5 form charge K carried by KK monopoles CMH

K-Charge in D=5

Spacetime M asymptotic to \bar{M}

k asymptotic to Killing vector on \bar{M}

$$\Delta\omega = \omega - \bar{\omega}$$

Difference in spin connections: Asymptotic tensor
ADM Momentum for k : Integral at spatial infinity Σ^3

$$P[k] = \frac{1}{16\pi^2} \int_{\Sigma^3} *(e_{\wedge}^A e_{\wedge}^B k)_{\wedge} \Delta\omega_{AB}$$

Nestor

K-charge

$$K = \frac{1}{16\pi^2} \int_{\Sigma^3} e_{\wedge}^A e_{\wedge}^B \Delta\omega_{AB}$$

Hull

K and NUT Charge

NUT Charge: Reduce on Killing vector
N is magnetic charge for graviphoton in D=4

KK Monopole spacetime: (Taub-NUT) \times (time)

NUT charge N

S^1 fibre, asymptotically radius $R=|N|$

$$K=RN=N|N|$$

Gravitational Instantons

Carry K

- $N \times (\text{time})$, N gravitational instanton
N Gibbons-Hawking multi-instanton space with general sources.
- Metric has Dirac string singularities in general, but connection well-defined
- If all charges are equal, singularities can be removed by identifying under discrete group: ALE or ALF instanton. But if not equal, singular.
- Should string singularities be allowed in quantum gravity? In M-theory?

Symmetry of (4,0)

Free theory:

Conventional field theory in flat background

Background diffeomorphisms + gauge trans

$$\delta C_{MNPQ} = \partial_{[M}\chi_{N]PQ} + \partial_{[P}\chi_{Q]MN} - 2\partial_{[M}\chi_{NPQ]}$$

Reduce to D=5 or D=4:

Combine $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$

2 Symmetries are the same for $g_{\mu\nu}$

On T^2 , background diffeos give $SL(2, \mathbb{Z})$ S-duality of both spin-1 and spin-2 fields in D=4

Interacting D=6 theory:

Can't combine background η_{MN} & field C_{MNPQ}

Don't expect D=6 diffeos, but exotic symmetries that give D=5 diffeomorphisms

Without D=6 diffeomorphisms, no reason to expect $SL(2, \mathbb{Z})$ and hence no "derivation" of gravitational S-duality (unlike free case)

Without D=6 diffeomorphisms, should spacetime be replaced by something more exotic?

This should be consistent with free limit being a conventional field theory

(2,0) & (4,0) 6-D CFTs

- No local covariant interacting field theory
- D=5 BPS electric 0-branes and magnetic strings lift to self-dual strings in D=6. Tension to zero in conformal limit
- Large superconformal symmetry: (4,0) has 32+32 susys
- YM and graviton in D=5 lift to self-dual tensor gauge fields
- D=5 g_{YM} & l_{planck} from R_6 as no scale in 6-D

M-Theory

- M-theory on T^6 has $D=5$ $N=8$ SUGRA as low energy limit
- $D=5$ branes lift to self-dual strings in $D=6$. Tension to zero in strong coupling limit
- Is strong coupling limit a 6D theory with $(4,0)$ SUSY, with exotic conformal gravity?
- Highly symmetric phase of M-theory?

Conclusions

- Dual gravitons and gravitational S-duality work well for free theory
- For $D \geq 5$, charge K carried by KK monopoles, and branes from $D=4$ instantons. Related to NUT charge and magnetic charge of KK monopoles
- For $D=4$ SYM or linearised SUGRA, S-duality from $(2,0)$ or $(4,0)$ theory on T^2

(4,0): All Four Nothing?

- Key question: are there BPS states with K?
- Extra dimension from strong coupling?
- (4,0) theory as a limit of M-theory?
Vast symmetry and unusual features
- Not usual spacetime, no metric or diffeos
- Is (4,0) CFT a decoupling limit of (4,0) sector of M-theory?

